arXiv:1010.3544v1 [math.NT] 18 Oct 2010

A Note on the Linnik's constant

Zaizhao MENG

1 Introduction

Suppose that q is a sufficiently large positive integer, $(a,q)=1$, let $P(a,q)$ be the least prime in the arithmetic progression $\{n\equiv a(mod\ q)\}$ and χ be the Dirichlet character of modulus q. $L(s,\chi)$ is the Dirichlet L-function. Every positive integer q can be expressed as $q=q_3^3q_2,\ q_2$ is cube-free. An integer q is called "has bounded cubic part" if the above q_3 is bounded absolutely.

In this note, we correct the Lemma of [4] and obtain the following result.

Theorem If q has bounded cubic part, for $(a,q)=1$, we have

$$P(a,q)\ll q^{4.5}.$$

Notation: ε is a sufficiently small positive real number. $\ell=\log q$, χ is the Dirichlet character of modulus q and χ_0 is the principal character, q is sufficiently large positive integer and has bounded cubic part.

2 Basic lemma and proof of the theorem

Let R be the rectangle (6.1) of [2]. Let ρ_1 be a zero of $\prod\limits_{\chi(mod\ q)}L(s,\chi)$ in R with $\beta_1=\Re(\rho_1)$ maximal, let χ_1 be any character with $L(\rho_1,\chi_1)=0$. Let ρ_2 be a zero of $\prod\limits_{\chi\neq\chi_1,\bar{\chi_1}}L(s,\chi)$ in R with $\beta_2=\Re(\rho_2)$ maximal, let χ_2 be any character with $L(\rho_2,\chi_2)=0$,.... We obtain new zeros ρ_1,..., and $\Re\rho_{k+1}\le\Re\rho_k\le...\le\Re\rho_2\le\Re\rho_1$. As in [2], we shall set$\rho_k=\beta_k+i\gamma_k,\ \beta_k=1-\ell^{-1}\lambda_k,\ \gamma_k=\ell^{-1}\mu_k$. The zero ρ' of $L(s,\chi)$ is as that in [2], $\rho'=\beta'+i\gamma',\ \beta'=1-\ell^{-1}\lambda',\ \gamma'=\ell^{-1}\mu'$.

Using the results in [1] and [2], set $\phi = \frac{1}{4}$ in the Lemma 3.1 of [2]. The bounds of $\lambda_1, \lambda_2, \lambda_3$ and λ' in [3] can be used to deduce the theorem. As in [3] and [4], we have $\lambda_3 > \frac{8}{7} - \varepsilon$ except for the case: $\chi_1\chi_2\chi_3 = \chi_0, \chi_1, \chi_2, \chi_3, \rho_1$ all are real.

Lemma. Suppose that $\chi_1\chi_2\chi_3 = \chi_0$, and $\chi_1, \chi_2, \chi_3, \rho_1$ all are real. If $\lambda_1 \leq 0.85$, then $\lambda_3 > \frac{8}{7} - \varepsilon$.

Proof. As in [4], suppose that $0.6 < \lambda_1 \leq 0.85$. Let $P_3(X) = X + X^2 + \frac{2}{3}X^3$, as in [2](on page 315), we can deduce that if $\lambda_3 \leq b \leq (\sqrt[3]{3} - 1)a$, then

$$P_3(\frac{a+\lambda_1}{a}) - P_3(1) - 2P_3(\frac{a+\lambda_1}{a+b}) + (a+\lambda_1)(\frac{3}{4} + \varepsilon) \geq 0. \tag{2.1}$$

We choose $b = 1.15,\ a = b(\sqrt[3]{3} - 1)^{-1}$, and will prove that when ε is sufficiently small and q is sufficiently large, (2.1) can not happen, and then $\lambda_3 > 1.15 > \frac{8}{7}$.

Let

$$S(X) = P_3(\frac{X}{a}) - P_3(1) - 2P_3(\frac{X}{a+b}) + \frac{3}{4}X,$$

then the differential function

$$S'(X) = c_1 + c_2X + c_3X^2,$$

where

$$c_1 = \frac{3}{4} + \frac{1}{a} - \frac{2}{a+b}, c_2 = \frac{2}{a^2} - \frac{4}{(a+b)^2}, c_3 = \frac{2}{a^3} - \frac{4}{(a+b)^3}.$$

$S'(X) = 0$ has not real root, and $S(a+0.6) = -0.269776...$, $S(a+0.85) = -0.0051...$, when ε is sufficiently small and q is sufficiently large, the left hand side of (2.1) is less than -0.005, this contradiction shows that $\lambda_3 > 1.15$. The lemma follows.

Suppose $N(\lambda)$ and H_0 have the same meaning as in [3]. Taking $\phi = \frac{1}{4}$ in Lemma 12.1 of [2], we add the following results.

$\lambda_1 = 0.85 : N(0.9) = 5,\ N(1.0) = 6,\ N(1.1) = 7,\ N(1.2) = 9,\ N(1.3) = 12,\ N(1.4) = 16,\ N(1.5) = 23$.

As in [3], we need to show $H_0 < 1$.

By the results of [3], it is necessary to add the following case , as in the proof of the Theorem 1.4 of [3]:

(a1) χ_1 is real, ρ_1 is real, $\lambda_1 \in [0.85, 1.15]$.

In this case (a1), using the above bounds for $N(\lambda)$, we deduce that $H_0 < 0.9$. This completes the proof of the theorem.

References

[1] D.A.Burgess, On character sums and L-series, II, Proc. London Math. Soc. (3) 13 (1963) 524-536.

[2] D.R.Heath-Brown, Zero-free regions for Dirichlet L-functions, and the least prime in an arithmetic progression, Proc. London Math. Soc. (3) 64 (1992) 265-338.

[3] Z.Z.Meng, The distribution of the zeros of L-functions and the least prime in some arithmetic progression, Sci. China Ser. A, 43 (2000) 937-944.

[4] Z.Z.Meng, A Note on the Least Prime in an Arithmetic Progression, Acta Sci. Natur. Univ. Pekinensis 37 (Jan 2001) 20-22.

School of Mathematical Sciences
Peking University
Beijing, 100871, P.R.China
E-mail: mzzh@math.pku.edu.cn